

Time-dependent CP Asymmetries in $B^0 \rightarrow K_S^0 \rho^0 \gamma$ Decays

J. Li,⁶ I. Adachi,⁷ K. Arinstein,¹ T. Aushev,^{17,12} A. M. Bakich,³⁷ V. Balagura,¹² I. Bedny,¹ V. Bhardwaj,³² U. Bitenc,¹³ A. Bozek,²⁶ M. Bračko,^{19,13} T. E. Browder,⁶ P. Chang,²⁵ Y. Chao,²⁵ A. Chen,²³ B. G. Cheon,⁵ R. Chistov,¹² Y. Choi,³⁶ J. Dalseno,⁷ A. Drutskoy,³ S. Eidelman,¹ N. Gabyshev,¹ H. Ha,¹⁵ K. Hara,²¹ Y. Hasegawa,³⁵ H. Hayashii,²² M. Hazumi,⁷ D. Heffernan,³¹ Y. Hoshi,⁴⁰ W.-S. Hou,²⁵ H. J. Hyun,¹⁶ T. Iijima,²¹ A. Ishikawa,³³ R. Itoh,⁷ M. Iwasaki,⁴² Y. Iwasaki,⁷ N. J. Joshi,³⁸ D. H. Kah,¹⁶ J. H. Kang,⁴⁶ H. Kawai,² T. Kawasaki,²⁸ H. Kichimi,⁷ Y. I. Kim,¹⁶ Y. J. Kim,⁴ K. Kinoshita,³ S. Korpar,^{19,13} P. Krizan,^{18,13} P. Krokovny,⁷ R. Kumar,³² A. Kuzmin,¹ S.-H. Kyeong,⁴⁶ C. Liu,³⁴ Y. Liu,⁴ A. Matyja,²⁶ S. McOnie,³⁷ T. Medvedeva,¹² K. Miyabayashi,²² H. Miyake,³¹ H. Miyata,²⁸ G. R. Moloney,²⁰ Y. Nagasaka,⁸ M. Nakao,⁷ Z. Natkaniec,²⁶ S. Nishida,⁷ O. Nitoh,⁴⁴ S. Ogawa,³⁹ T. Ohshima,²¹ S. Okuno,¹⁴ S. L. Olsen,^{6,9} H. Ozaki,⁷ G. Pakhlova,¹² C. W. Park,³⁶ H. Park,¹⁶ H. K. Park,¹⁶ K. S. Park,³⁶ L. S. Peak,³⁷ R. Pestotnik,¹³ L. E. Piilonen,⁴⁵ H. Sahoo,⁶ Y. Sakai,⁷ O. Schneider,¹⁷ C. Schwanda,¹⁰ A. Sekiya,²² K. Senyo,²¹ M. Shapkin,¹¹ J.-G. Shiu,²⁵ B. Shwartz,¹ A. Sokolov,¹¹ A. Somov,³ S. Stanič,²⁹ M. Starič,¹³ T. Sumiyoshi,⁴³ M. Tanaka,⁷ G. N. Taylor,²⁰ Y. Teramoto,³⁰ I. Tikhomirov,¹² K. Trabelsi,⁷ T. Tsuboyama,⁷ S. Uehara,⁷ T. Uglov,¹² Y. Unno,⁵ S. Uno,⁷ P. Urquijo,²⁰ Y. Ushiroda,⁷ Y. Usov,¹ G. Varner,⁶ K. E. Varvell,³⁷ K. Vervink,¹⁷ A. Vinokurova,¹ C. H. Wang,²⁴ P. Wang,⁹ X. L. Wang,⁹ Y. Watanabe,¹⁴ E. Won,¹⁵ B. D. Yabsley,³⁷ H. Yamamoto,⁴¹ Y. Yamashita,²⁷ M. Yamauchi,⁷ Z. P. Zhang,³⁴ V. Zhilich,¹ T. Zivko,¹³ A. Zupanc,¹³ and O. Zyukova¹

(The Belle Collaboration)

¹*Budker Institute of Nuclear Physics, Novosibirsk*

²*Chiba University, Chiba*

³*University of Cincinnati, Cincinnati, Ohio 45221*

⁴*The Graduate University for Advanced Studies, Hayama*

⁵*Hanyang University, Seoul*

⁶*University of Hawaii, Honolulu, Hawaii 96822*

⁷*High Energy Accelerator Research Organization (KEK), Tsukuba*

⁸*Hiroshima Institute of Technology, Hiroshima*

⁹*Institute of High Energy Physics, Chinese Academy of Sciences, Beijing*

¹⁰*Institute of High Energy Physics, Vienna*

¹¹*Institute of High Energy Physics, Protvino*

¹²*Institute for Theoretical and Experimental Physics, Moscow*

¹³*J. Stefan Institute, Ljubljana*

¹⁴*Kanagawa University, Yokohama*

¹⁵*Korea University, Seoul*

¹⁶*Kyungpook National University, Taegu*

¹⁷*École Polytechnique Fédérale de Lausanne (EPFL), Lausanne*

¹⁸*Faculty of Mathematics and Physics, University of Ljubljana, Ljubljana*

¹⁹*University of Maribor, Maribor*

²⁰*University of Melbourne, School of Physics, Victoria 3010*

²¹*Nagoya University, Nagoya*

²²*Nara Women's University, Nara*

²³*National Central University, Chung-li*

²⁴*National United University, Miao Li*

²⁵*Department of Physics, National Taiwan University, Taipei*

²⁶*H. Niewodniczanski Institute of Nuclear Physics, Krakow*

²⁷*Nippon Dental University, Niigata*

²⁸*Niigata University, Niigata*

²⁹*University of Nova Gorica, Nova Gorica*

³⁰*Osaka City University, Osaka*

³¹*Osaka University, Osaka*

³²*Panjab University, Chandigarh*

³³*Saga University, Saga*

³⁴*University of Science and Technology of China, Hefei*

³⁵*Shinshu University, Nagano*

³⁶*Sungkyunkwan University, Suwon*

³⁷*University of Sydney, Sydney, New South Wales*

³⁸Tata Institute of Fundamental Research, Mumbai

³⁹Toho University, Funabashi

⁴⁰Tohoku Gakuin University, Tagajo

⁴¹Tohoku University, Sendai

⁴²Department of Physics, University of Tokyo, Tokyo

⁴³Tokyo Metropolitan University, Tokyo

⁴⁴Tokyo University of Agriculture and Technology, Tokyo

⁴⁵Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061

⁴⁶Yonsei University, Seoul

We report the first measurement of CP -violation parameters in $B^0 \rightarrow K_S^0 \rho^0 \gamma$ decays based on 657 million $B\bar{B}$ pairs collected with the Belle detector at the KEKB asymmetric-energy collider. We measure the time-dependent CP violating parameter $\mathcal{S}_{K_S^0 \rho^0 \gamma} = 0.11 \pm 0.33(\text{stat.})_{-0.09}^{+0.05}(\text{syst.})$. We also obtain the effective direct CP violating parameter $\mathcal{A}_{\text{eff}} = 0.05 \pm 0.18(\text{stat.}) \pm 0.06(\text{syst.})$ for $m_{K_S \pi^+ \pi^-} < 1.8 \text{ GeV}/c^2$ and $0.6 \text{ GeV}/c^2 < m_{\pi^+ \pi^-} < 0.9 \text{ GeV}/c^2$.

PACS numbers: 11.30.Er, 13.20.He

In the standard model (SM), a mostly left (right)-handed photon emitted from a $\bar{B}^0$ (B^0) meson is expected in the $b \rightarrow s\gamma$ transition. Hence a small time-dependent CP asymmetry is predicted in decays of the type $B \rightarrow f_{CP}\gamma$ [1], where f_{CP} is a CP eigenstate. New Physics (NP) may lead to deviations from the SM expectation by introducing different photon polarizations in the transition, and can be probed via experimental measurements of CP violation parameters [2]. In multi-body final states $B^0 \rightarrow P^0 Q^0 \gamma$ with P^0 and Q^0 being C eigenstates, the same argument holds [3]. Measurements of the time-dependent CP asymmetry in $B^0 \rightarrow K_S^0 \pi^0 \gamma$ have been reported by Belle and BaBar based on 535 and 431 million $B\bar{B}$ pairs [4, 5], respectively. In this paper, based on 657 million $B\bar{B}$ pairs collected with the Belle detector [6] at the KEKB asymmetric-energy e^+e^- collider [7], we report the measurement of CP -violation parameters on a new channel $B^0 \rightarrow K_S^0 \rho^0 \gamma$ where the B^0 decay vertex can be reconstructed from two charged pions from the ρ^0 decays.

At the KEKB, the $\Upsilon(4S)$ is produced with a Lorentz boost of $\beta\gamma = 0.425$ along the z axis, which is defined as the direction antiparallel to the e^+ beam direction. In the decay chain $\Upsilon(4S) \rightarrow B^0 \bar{B}^0 \rightarrow f_{\text{sig}} f_{\text{tag}}$, where one of the B mesons decays at time t_{sig} to the signal mode f_{sig} and the other decays at time t_{tag} to a final state f_{tag} that distinguishes B^0 and $\bar{B}^0$, the time-dependent decay rate is given by:

$$P(\Delta t, q) = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} \left\{ 1 + q \cdot \left[\mathcal{S} \sin(\Delta m_d \Delta t) + \mathcal{A} \cos(\Delta m_d \Delta t) \right] \right\}. \quad (1)$$

Here the probability density function (PDF) is normalized as a function of two variables: B flavor $q = +1(-1)$ when the tagging B meson is $B^0(\bar{B}^0)$, and decay time difference Δt between two B mesons. In Eq.(1), $\mathcal{S}$ and $\mathcal{A}$ are CP -violation parameters, τ_{B^0} is the B^0 lifetime,

Δm_d is the mass difference between the two B^0 mass eigenstates. Since the $B^0 \bar{B}^0$ mesons are approximately at rest in the $\Upsilon(4S)$ center-of-mass system (c.m.s.), Δt can be determined from Δz , the displacement in z between the f_{sig} and f_{tag} decay vertices: $\Delta t \simeq \Delta z/(\beta\gamma c)$.

The Belle detector [6] is a large-solid-angle magnetic spectrometer that consists of a silicon vertex detector (SVD), a 50-layer central drift chamber, an array of aerogel threshold Cherenkov counters, and an electromagnetic calorimeter (ECL) comprised of CsI(Tl) crystals located inside a superconducting solenoid coil that provides a 1.5 T magnetic field. An iron flux-return located outside the coil is instrumented to detect K_L^0 mesons and identify muons.

High energy photons are selected from isolated ECL clusters with no corresponding charged track and c.m.s. energy satisfying $1.4 \text{ GeV} < E_\gamma^* < 3.4 \text{ GeV}$. The photons are also required to lie in the barrel region of the ECL, and have a photon shower-like shape $E_9/E_{25} > 0.95$, where E_9 and E_{25} are the energies summed in 3×3 and 5×5 arrays of crystals around the center of the shower, respectively. To reduce background from $\pi^0 \rightarrow \gamma\gamma$ or $\eta \rightarrow \gamma\gamma$, a $\pi^0(\eta)$ veto is applied with $\mathcal{L}_{\pi^0} < 0.25$ ($\mathcal{L}_\eta < 0.2$), where $\mathcal{L}_{\pi^0(\eta)}$ is a $\pi^0(\eta)$ likelihood described in Ref. [8].

Neutral kaons (K_S^0) are reconstructed from two oppositely charged pions whose invariant mass lies within 15 MeV/c^2 of the K_S^0 nominal mass. Requirements on impact parameter and vertex displacement are applied [9].

Charged tracks are required to originate from the vicinity of the interaction point (IP). Charged pions should have kaon and pion identification likelihoods consistent with the pion hypothesis. This requirement has an efficiency of 85% with a 7% kaon fake rate.

The decay $B^+ \rightarrow K^+ \pi^- \pi^+ \gamma$ is also reconstructed to study the $K\pi\pi$ system and to serve as a control sample. Charged kaons are selected from charged tracks and required to be identified as a kaon with 86% efficiency and an 8% pion fake rate.

We form two kinematic variables: the energy difference

$\Delta E = E_B^* - E_{\text{beam}}^*$ and the beam-energy constrained mass $M_{bc} = \sqrt{(E_{\text{beam}}^*)^2 - (p_B^*)^2}$, where E_{beam}^* is the beam energy in the c.m.s., and E_B^* and p_B^* are the c.m.s. energy and momentum of the reconstructed B candidate, respectively. The requirement $-0.1 \text{ GeV} < \Delta E < 0.08 \text{ GeV}$ is applied. The $K^+\pi^-\pi^+$ and $K_S^0\pi^-\pi^+$ invariant masses are required to be less than $1.8 \text{ GeV}/c^2$.

The $B^0 \rightarrow K_S^0\rho^0\gamma$ candidates are selected from the $K_S^0\pi^+\pi^-\gamma$ sample by requiring the $\pi^+\pi^-$ invariant mass to lie in the ρ^0 region, $0.6 \text{ GeV}/c^2 < m_{\pi\pi} < 0.9 \text{ GeV}/c^2$. Since the ρ^0 is wide, other modes, such as $K^{*+}\pi^-\gamma$ may also contribute. We first measure the effective CP -violating parameters, $\mathcal{S}_{\text{eff}}$ and $\mathcal{A}_{\text{eff}}$, using the final sample and then convert them to the CP -violating parameters of $B^0 \rightarrow K_S^0\rho^0\gamma$ using a dilution factor $\mathcal{D}$, which is discussed later.

In order to suppress the background from light quark pair production $q\bar{q}$ ($e^+e^- \rightarrow q\bar{q}$ with $q = u, d, s, c$), the selection based on an event likelihood ratio $\mathcal{R} \equiv \mathcal{L}_{\text{sig}}/(\mathcal{L}_{\text{sig}} + \mathcal{L}_{\text{bkg}})$ is applied. The likelihood for signal ($\mathcal{L}_{\text{sig}}$) and background ($\mathcal{L}_{\text{bkg}}$) is formed by combining a Fisher discriminant $\mathcal{F}$ that uses extended modified Fox-Wolfman moments [10] and the polar angle of the B meson in the c.m.s. ($\cos\theta_B$). If there are multiple candidates in an event, we choose the candidate that has the largest $\mathcal{R}$. We define ΔE sideband events with the criteria $0.1 \text{ GeV} < \Delta E < 0.5 \text{ GeV}$ and by vetoing $B \rightarrow (K\pi^\pm)\gamma$. Other backgrounds that pass our selection criteria are classified as (a) $B \rightarrow K^*\gamma$ background, (b) $B \rightarrow X_s\gamma$ background, (c) $B\bar{B}$ background, which includes a generic $b \rightarrow c$ component and charmless B -decay component.

The self-cross-feed (SCF) component consists of events in which not all tracks are from the signal side, in contrast to a true signal with all tracks correctly assigned. We define the total signal yield as the sum of SCF and true signal events. The fraction of SCF in the total signal yield ranges from 5.5% to 10.8% in the M_{bc} signal region defined as $5.27 \text{ GeV}/c^2 < M_{bc} < 5.29 \text{ GeV}/c^2$, depending on the kaonic resonance.

We obtain the flavor q , tagging quality factor $r \in [0, 1]$, and signal and tag side vertices from the procedure described in Ref. [11], using the two charged pions to reconstruct the signal vertex position. There is no flavor discrimination when $r = 0$, while the flavor tagging is unambiguous when $r = 1$. Events are sorted into seven r -bins.

We obtain 299 events in the signal M_{bc} region after vertexing. An unbinned maximum likelihood fit to M_{bc} is applied with signal and $q\bar{q}$ yields floated in each r -bin. In the signal M_{bc} region, we find 212 ± 17 total signal with a fraction of 6.0% SCF, 53.4 ± 2.6 continuum, along with 7.8 $K^*\gamma$, 21.5 other $X_s\gamma$, and 9.0 $B\bar{B}$ events. The true signal shape is parameterized as a Crystal Ball function with a width corrected using the $B^+ \rightarrow K^+\pi^-\pi^+\gamma$ data. The $q\bar{q}$ shape is obtained from the ΔE sideband events.




FIG. 1: M_{bc} distributions for $B^0 \rightarrow K_S^0\pi^+\pi^-\gamma$ events. Points with error bars are data. The curves show the results from the r dependent M_{bc} fit. The dashed and dash-dotted curves are the $q\bar{q}$ and all BG. The thin curve is the total signal including SCF and the thick curve is the total PDF.

For other backgrounds and SCF, shapes and r -dependent fractions are obtained from MC since their contributions are limited. Figure 1 shows the M_{bc} projection of the fit.

The $\mathcal{S}_{\text{eff}}$ and $\mathcal{A}_{\text{eff}}$ parameters are extracted from an unbinned maximum likelihood fit to the Δt distribution. The likelihood function for each event is:

$$P_i = (1 - f_{\text{ol}}) \int_{-\infty}^{+\infty} d(\Delta t') \left[\sum_j f_j P_j(\Delta t') R_j(\Delta t_i - \Delta t') \right] + f_{\text{ol}} P_{\text{ol}}(\Delta t_i), \quad (2)$$

where j runs over the signal, SCF and four BG components ($q\bar{q}$, $K^*\gamma$, other $X_s\gamma$, $B\bar{B}$). The fraction of each component (f_j) is calculated using the r dependent M_{bc} fit result on an event-by-event basis. R_j is the resolution function in the Δt .

The PDF for the signal Δt distribution P_{sig} , is given by a modified form of Eq.(1), which incorporates the effect of incorrect flavor assignment. The parameterization of R is the same as the one used in the $B^0 \rightarrow \phi K^0$ analysis [9]. The PDF for SCF (P_{SCF}) is the same as for the signal except for using a shorter lifetime $1.16 \pm 0.02 \text{ ps}$, determined from the MC study. The same functional forms for the PDF and resolution are used for the $K^*\gamma$, other $X_s\gamma$ and $B\bar{B}$ components but with other lifetime values obtained from MC and CP parameters fixed to zero. The PDF for $q\bar{q}$ background events, $P_{q\bar{q}}$, is modeled as a δ function convolved with a double-Gaussian resolution function $R_{q\bar{q}}$. The parameters in $R_{q\bar{q}}$ is determined from a fit to ΔE sideband events. P_{ol} is a Gaussian function that represents a small outlier component with a fraction f_{ol} .

The only free parameters in the CP fit to $B^0 \rightarrow K_S^0\pi^+\pi^-\gamma$ are $\mathcal{S}_{\text{eff}}$ and $\mathcal{A}_{\text{eff}}$, which are determined by maximizing the likelihood function $L = \prod_i P_i$ for events in the M_{bc} signal region. We obtain $\mathcal{S}_{\text{eff}} = 0.09 \pm 0.27(\text{stat.})^{+0.04}_{-0.07}(\text{syst.})$ and $\mathcal{A}_{\text{eff}} = 0.05 \pm 0.18(\text{stat.}) \pm$




FIG. 2: Fit projections on the Δt distributions with (a) $q = +1$ and (b) $q = -1$ for events with $r > 0.5$. The solid curves are the fit while the dashed curves show the background contributions. The raw asymmetry as a function of Δt is shown in (c) with a fit curve superimposed.

0.06(syst.). We define the raw asymmetry in each Δt bin by $(N_+ - N_-)/(N_+ + N_-)$, where N_+ (N_-) is the number of candidate events with $q = +1$ (-1). Figure 2 shows the Δt distributions and the raw asymmetry for events with $0.5 < r \leq 1.0$.

Various validity checks for our fitting procedure are performed. Lifetime fit results for the $B^+ \rightarrow K^+\pi^-\pi^+\gamma$ and $B^0 \rightarrow K_S^0\pi^+\pi^-\gamma$ modes are consistent with the nominal B^+ and B^0 lifetimes. No CP asymmetry is seen for the control sample $B^+ \rightarrow K^+\pi^-\pi^+\gamma$.

We evaluate systematic uncertainties from the following sources. The largest contribution is due to the vertex reconstruction, where the selection criteria are varied to calculate the systematics as $\Delta\mathcal{S}_{\text{eff}} = {}^{+0.01}_{-0.06}$, $\Delta\mathcal{A}_{\text{eff}} = \pm 0.03$. The $\mathcal{S}$ values for $b \rightarrow s$ CP modes are varied from 0 to $\sin 2\phi_1$ to estimate the effect of CP asymmetry in the $B\bar{B}$ background. We fit the data with each fixed parameter shifted by its error to evaluate the uncertainties due to: background and SCF fractions, shapes, Δt PDFs, resolution function, flavor tagging and physics parameters τ_{B^0} , Δm_d . Effects of tag side interference are evaluated in the same way as in Ref. [4].

The parameter $\mathcal{S}_{\text{eff}}$ is related to $\mathcal{S}$ for $K_S^0\rho^0\gamma$ with a dilution factor $\mathcal{D}$ that depends on the $K^{*\pm}\pi^\mp$ components:

$$\mathcal{D} = \frac{\mathcal{S}_{\text{eff}}}{\mathcal{S}_{K_S^0\rho^0\gamma}} = \frac{\int [|F_A|^2 + 2\Re(F_A^*F_B) + F_B^*(\bar{K})F_B(K)]}{\int [|F_A|^2 + 2\Re(F_A^*F_B) + |F_B|^2]}, \quad (3)$$

where F_A, F_B are photon-helicity averaged amplitudes for $B^0 \rightarrow K_S^0\rho^0(\pi^+\pi^-)\gamma$ and $B^0 \rightarrow K^{*\pm}(K_S^0\pi^\pm)\pi^\mp\gamma$, respectively. The factors $F_B(\bar{K})$, $F_B(K)$ distinguish between $K^{*-}\pi^+\gamma$ and $K^{*+}\pi^-\gamma$. The phase space integral is over the ρ^0 region.

The charged mode $B^+ \rightarrow K^+\pi^-\pi^+\gamma$ is first studied using a combination of various kaonic resonances with spin ≥ 1 to model the $K\pi\pi$ system. The amplitude for a kaonic resonance K_{res} that decays into particle c and

subresonance r , where r later decays into particles a and b , can be modeled by the product of two Breit-Wigners:

$$M_{abc|r} = BW(K_{\text{res}})BW(r)F_K(K_{\text{res}})F_r f_{\text{spin}}, \quad (4)$$

where BW is the relativistic Breit-Wigner lineshape: $BW(K_{\text{res}}) = 1/(m_{K_{\text{res}}}^2 - m_{abc}^2 - im_{K_{\text{res}}}\Gamma_{K_{\text{res}}})$ and $BW(r) = 1/(m_r^2 - m_{ab}^2 - im_r\Gamma_{ab})$. The width $\Gamma_{ab} = \Gamma_r(q/q_0)^{2L+1} \left(\frac{m_r}{m_{ab}}\right) F_r^2$ is a function of q and q_0 , the momenta of particle a in the r rest frame with mass m_{ab} and m_r , respectively. $F_K(K_{\text{res}})$ and F_r are the Blatt-Weisskopf penetration factors [12] for resonances K_{res} and r . The spin factor f_{spin} for the resonances used in the nominal fit is: $f_{\text{spin}} = 1$ for $K_1^+(1270) \rightarrow K\rho^0, K^*\pi$, $K_1^+(1400) \rightarrow K^*\pi$ with a 1^+S wave; $f_{\text{spin}} = \sin\theta$ for $K^*(1680) \rightarrow K\rho^0, K^*\pi$ with a 1^-P wave; $f_{\text{spin}} = \sin\theta$ for $K_2^+(1430) \rightarrow K^*\pi, K\rho^0$ with a 2^+D wave. Here θ is the helicity angle of subresonance r .

Kaonic resonances with different spin-parity waves do not interfere if the decay plane orientation variables are integrated out. The total rate is an incoherent sum of contributions from different spin-parities with a phase space factor.

To determine the $K^+\rho^0\gamma$ component in $B^+ \rightarrow K^+\pi^+\pi^-\gamma$, we study events in a K^* region defined as $|m_{K^+\pi^-} - 0.8961| < 0.075$ GeV/ c^2 , where most of the $B^+ \rightarrow K^+\pi^+\pi^-\gamma$ signal lies. We fit the $m_{K\pi\pi}, m_{\pi^+\pi^-}, m_{K^+\pi^-}$ distributions for events in the signal M_{bc} region. In these fits, the yields and shapes for backgrounds are obtained from the corresponding M_{bc} fit results and MC. The fraction and shape for SCF are associated with each kaonic resonance from MC. The yields for the $2^+ K_2^*(1430)\gamma$ components are always fixed based on the measured branching fraction $\mathcal{B}(B \rightarrow K_2^*(1430)\gamma) = (1.24 \pm 0.24) \times 10^{-5}$ [13].

A two-dimensional (2D) $m_{K\pi\pi}, m_{\pi\pi}$ fit is performed in the K^* region. Here the SCF is further categorized depending on whether the K^+, π^- tracks are correctly reconstructed. The floating parameters are the rate and phase for $K_1(1270) \rightarrow K^*\pi$ relative to $K_1(1270) \rightarrow K\rho$. In the fit, the yields of $1^+, 1^-$ kaonic resonances are obtained from a fit to the $m_{K\pi\pi}$ distribution in the K^* region. The rates for $K^+\rho^0$ relative to $K^{*0}\pi^+$ in the $K^*(1680)$ and $K_2^*(1430)$ resonances are fixed according to the PDG values [13], taking into account phase-space and isospin factors. The phases for $K_{\text{res}} \rightarrow K^{*0}\pi^+$ relative to $K_{\text{res}} \rightarrow K^+\rho^0$ in the $K^*(1680)$, $K_2^*(1430)$ resonances are fixed to be -0.36 rad [14] and -30° [15], respectively. The PDF for the $m_{K\pi\pi}$ distribution in the ρ^0 region is then obtained.

The procedure is repeated for different fixed values of the phase ϕ for $K_1(1400) \rightarrow K^*\pi$ relative to $K_1(1270) \rightarrow K\rho$. For the nominal fit, we use $\phi = 15^\circ$ which gives the smallest total χ^2 for the 2D $m_{K\pi\pi}, m_{\pi\pi}$ distribution in the K^* region and $m_{K\pi\pi}$ distribution in the ρ^0 region. Figure 3(a) shows the distributions for $m_{\pi\pi}$ in the whole




FIG. 3: $m_{\pi\pi}$ distributions for (a) $B^+ \rightarrow K^+\pi^-\pi^+\gamma$ and (b) $B^0 \rightarrow K_S^0\pi^+\pi^-\gamma$. The curves follow the convention in Fig. 1. The thin dashed curve is the correctly reconstructed $B \rightarrow K_1(1270)\gamma$ signal.

TABLE I: The number of events for each kaonic resonance and various interference terms for the final state $K_S^0\pi^+\pi^-\gamma$. Interf. denotes the interference between $K_S^0\rho^0\gamma$ and $K^{*+}\pi^-\gamma = 2\Re(F_A^*F_B)$.

	Total	$K_S^0\rho^0\gamma$	$K^{*+}\pi^-\gamma$	Interf.	$\frac{F_B^*(\bar{K})}{F_B(K)}$
$K_{\text{res}}(1^+)\gamma$	193.6	151.0	35.1	7.5	4.4
$(K_1^0(1270)\gamma)$	(167.6)	(151.0)	(38.0)	(-21.4)	(5.2)
$K_{\text{res}}(1^-)\gamma$	24.2	11.3	8.0	4.9	1.3
$K_2^{*0}(1430)\gamma$	10.4	2.2	6.1	2.0	4.5
Sum	228.1	164.4	49.2	14.5	10.2

region. The data distributions for $m_{K\pi\pi}, m_{K\pi}, m_{\pi\pi}$ in other regions also agree with our model.

Using isospin symmetry, we assume that the fraction and phase of each kaonic resonance channel in the B^0 decay is the same as that in the B^+ decay. Therefore, using the fit results for the kaonic resonance composition and substituting the neutral kaonic resonances and K_S^0 mass instead of K^+ mass, we obtain the contributions of the terms $|F_A|^2, |F_B|^2$ in Eq.(3), as listed in Table I. Figure 3(b) shows the $m_{\pi\pi}$ distributions for $B^0 \rightarrow K_S^0\pi^+\pi^-\gamma$.

Various systematic uncertainties in the dilution factor have been investigated. The dominant one comes from the modeling systematics. We put additional resonances to the default model and repeated the fit procedure. For $B \rightarrow K\sigma\gamma$ in which the σ has a C parity opposite to that of the ρ^0 , the dilution factor is:

$$\mathcal{D} = \frac{\int [F_-(K)^*F_-(\bar{K}) - F_+^*F_+]}{\int |F_- + F_+|^2}. \quad (5)$$

Here F_+ is the photon-helicity averaged amplitude for $K_1(1270)(\rightarrow K_S^0\sigma)\gamma$, while $F_- = F_A + F_B$ [Eq.(3)]. In the absence of F_+ , Eq.(5) reduces to Eq.(3). The $\mathcal{D}$ value in this case is calculated to be 1.00. Other systematics con-

sidered include: model uncertainty from $K^{*0}(1430)\pi\gamma$, $K_1(1400)\gamma$, $K_1(1270) \rightarrow K\omega$ components, uncertainty in subresonance rates and phases, SCF uncertainty, isospin breaking, the mass and width of kaonic resonances, and statistical uncertainty from data. Adding all the uncertainties in quadrature, we obtain $\mathcal{D} = 0.83^{+0.19}_{-0.03}$.

By combining $\mathcal{S}_{\text{eff}}$ and the dilution factor $\mathcal{D}$, we obtain $\mathcal{S}_{K_S\rho^0\gamma} = 0.11 \pm 0.33(\text{stat.})^{+0.05}_{-0.09}(\text{syst.})$.

In summary, we have measured the time-dependent CP asymmetry in the decay $B^0 \rightarrow K_S\rho^0\gamma$ using events with $m_{K\pi\pi} < 1.8 \text{ GeV}/c^2$ and $0.6 \text{ GeV}/c^2 < m_{\pi\pi} < 0.9 \text{ GeV}/c^2$. We obtain CP -violation parameters $\mathcal{S}_{K_S\rho^0\gamma} = 0.11 \pm 0.33(\text{stat.})^{+0.05}_{-0.09}(\text{syst.})$ and $\mathcal{A}_{\text{eff}} = 0.05 \pm 0.18(\text{stat.}) \pm 0.06(\text{syst.})$. With the present statistics, the result is consistent with zero and comparable in precision to the measurement in $B^0 \rightarrow K_S^0\pi^0\gamma$. This is the first measurement of CP asymmetry parameters in the $K_S\rho^0\gamma$ mode and constrains right-handed currents from NP.

We thank the KEKB group for excellent operation of the accelerator, the KEK cryogenics group for efficient solenoid operations, and the KEK computer group and the NII for valuable computing and SINET3 network support. We acknowledge support from MEXT and JSPS (Japan); ARC and DEST (Australia); NSFC (China); DST (India); MOEHRD, KOSEF and KRF (Korea); KBN (Poland); MES and RFAAE (Russia); ARRS (Slovenia); SNSF (Switzerland); NSC and MOE (Taiwan); and DOE (USA).

- [1] D. Atwood, M. Gronau and A. Soni, Phys. Rev. Lett. **79**, 185 (1997) [arXiv:hep-ph/9704272].
- [2] T. Gershon and A. Soni, J. Phys. G **33**, 479 (2007) [arXiv:hep-ph/0607230].
- [3] D. Atwood, T. Gershon, M. Hazumi and A. Soni, Phys. Rev. D **71**, 076003 (2005) [arXiv:hep-ph/0410036].
- [4] Y. Ushiroda *et al.* [Belle Collaboration], Phys. Rev. D **74**, 111104 (2006) [arXiv:hep-ex/0608017].
- [5] B. Aubert *et al.* [BaBar Collaboration], arXiv:0708.1614 [hep-ex].
- [6] A. Abashian *et al.* [Belle Collaboration], Nucl. Instrum. Meth. A **479**, 117 (2002).
- [7] S. Kurokawa and E. Kikutani, Nucl. Instr. and. Meth. A **499**, 1 (2003), and other papers included in this volume.
- [8] P. Koppenburg *et al.* [Belle Collaboration], Phys. Rev. Lett. **93**, 061803 (2004) [arXiv:hep-ex/0403004].
- [9] K. F. Chen *et al.* [Belle Collaboration], Phys. Rev. Lett. **98**, 031802 (2007) [arXiv:hep-ex/0608039].
- [10] S. H. Lee *et al.* [Belle Collaboration], Phys. Rev. Lett. **91**, 261801 (2003) [arXiv:hep-ex/0308040].
- [11] Y. Ushiroda *et al.* [Belle Collaboration], Phys. Rev. Lett. **100**, 021602 (2008) [arXiv:0709.2769 [hep-ex]].
- [12] J. Blatt and V. Weisskopf, *Theoretical Nuclear Physics*, New York: John Wiley & Sons (1952). The penetration factor is defined as 1 for S wave, $\frac{\sqrt{R^2 p^2}}{\sqrt{1+R^2 p^2}}$ for P wave,

and $\sqrt{\frac{(Rp)^4}{(Rp)^4+3(Rp)^2+9}}$ for D wave, where p denotes the momentum of the daughter particle in the rest frame of the mother particle. We choose the radius to be $R = 0.3$ fm.

[13] W. M. Yao *et al.* [Particle Data Group], J. Phys. G **33**,

1 (2006).

[14] D. Aston *et al.*, Nucl. Phys. B **292**, 693 (1987).

[15] C. Daum *et al.* [ACCMOR Collaboration], Nucl. Phys. B **187**, 1 (1981).